
# CXXC4 Antibody (F55032)


| Catalog No. | Formulation | Size |
|---|---|---|
| F55032-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F55032-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

**Bulk quote request**


| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human, Mouse |
| **Format** | Purified |
| **Host** | Rabbit |
| **Clonality** | Polyclonal (rabbit origin) |
| **Isotype** | Rabbit Ig |
| **Purity** | Antigen affinity purified |
| **UniProt** | Q9H2H0 |
| **Localization** | Cytoplasmic |
| **Applications** | Western Blot : 1:500-1:1000<br>Flow Cytometry : 1:10-50 (1x10e6 cells)<br>Immunohistochemistry (FFPE) : 1:50-1:100 |
| **Limitations** | This CXXC4 antibody is available for research use only. |





Western blot testing of human K562 cell lysate with CXXC4 antibody. Predicted molecular weight ~21 kDa.




Western blot testing of mouse testis tissue lysate with CXXC4 antibody. Predicted molecular weight ~21 kDa.



IHC testing of FFPE human colon carcinoma tissue with CXXC4 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.




Flow cytometry testing of human K562 cells with CXXC4 antibody; Blue=isotype control, Green= CXXC4 antibody.

## Description

CXXC4 acts as a negative regulator of the Wnt signaling pathway via its interaction with DVL1.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the CXXC4 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 158-185 from the human protein was used as the immunogen for the CXXC4 antibody.

## Storage

Aliquot the CXXC4 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.