


CYP3A1 Antibody [clone P6] (V8294)

Catalog No.	Formulation	Size
V8294-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8294-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8294SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	P6
Purity	Protein G affinity chromatography
UniProt	P04800
Applications	ELISA (order BSA-free Format For Coating) : Flow Cytometry : 1-2ug/10 ⁶ cells Western Blot : 1-2ug/ml
Limitations	This CYP3A1 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free CYP3A1 antibody (clone P6) as confirmation of integrity and purity.

Description

Cytochrome P450 oxidase (commonly abbreviated CYP) is a generic term for a large number of related, but distinct, oxidative enzymes important in vertebrate physiology. The cytochrome P450 mixed-function monooxygenase system is

probably the most important element of Phase I metabolism in mammals. P450s are membrane-bound, either in the inner membrane of mitochondria or in the endoplasmic reticulum of cells where they metabolise thousands of endogenous and exogenous compounds. In the liver, these substrates include toxins, drugs, and other unneeded and potentially harmful molecules. Humans have 18 families of cytochrome P450 genes and 43 subfamilies; the CYP1 family is involved in drug metabolism and includes 3 subfamilies, 3 genes and 1 pseudogene. The CYP2 family are part of the microsomal drug metabolising system that is responsible for oxidation of many therapeutic agents as well as steroids, fatty acids and many other endogenous substances. Cytochrome P450 3A4 is abundantly expressed in liver and small intestine and is inducible by barbiturates, glucocorticoids and rifampicin. P6 is highly specific for members of the CYP 3A gene family and cross reacts with CY 3A4 in human, CYP 3A in rat and potentially CYP 3A11 (based on inducibility) in mice.

Application Notes

Optimal dilution of the CYP3A1 antibody should be determined by the researcher.

Immunogen

Rat liver cytochrome P450 3A1 was used as the immunogen for the CYP3A1 antibody.

Storage

Store the CYP3A1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).