
Cytokeratin 7 Antibody KRT7/760 / Mouse Monoclonal [clone KRT7/760] (V2658)

Catalog No.	Formulation	Size
V2658-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2658-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2658SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2658IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	KRT7/760
Purity	Protein G affinity chromatography
UniProt	P08729
Localization	Cytoplasmic
Applications	Flow Cytometry : 1-2ug/10^6 cells Immunofluorescence : 1-2ug/ml Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Cytokeratin 7 Antibody KRT7/760 is available for research use only.




Cytokeratin 7 Antibody KRT7/760 IHC: Formalin-fixed, paraffin-embedded human lung carcinoma stained with Cytokeratin 7 antibody (clone KRT7/760).




Cytokeratin 7 Antibody KRT7/760 IHC: Formalin-fixed, paraffin-embedded human ovarian carcinoma stained with Cytokeratin 7 antibody (clone KRT7/760).




Cytokeratin 7 Antibody KRT7/760 IHC: Formalin-fixed, paraffin-embedded human endometrial carcinoma stained with Cytokeratin 7 antibody (clone KRT7/760).



Cytokeratin 7 Antibody KRT7/760 WB Analysis. Western blot testing of human HeLa cell lysate with Cytokeratin 7 antibody (clone KRT7/760). Predicted molecular weight ~51 kDa.



Cytokeratin 7 Antibody KRT7/760 FACS Analysis. Flow cytometry testing of PFA-fixed human HeLa cells with Cytokeratin 7 antibody (clone KRT7/760); Red=isotype control, Blue= recombinant Cytokeratin 7 antibody.



Cytokeratin 7 Antibody KRT7/760 IF Analysis. Immunofluorescent staining of methanol-fixed human HeLa cells with Cytokeratin 7 antibody (clone KRT7/760, green) and Reddot nuclear stain (red).

Description

Cytokeratin 7 Antibody KRT7/760 is a monoclonal antibody specific for keratin 7, a type II intermediate filament protein expressed in many glandular and transitional epithelia. Keratin 7 is absent in gastrointestinal epithelia, making it a valuable marker for differential diagnosis of carcinomas. NSJ Bioreagents supplies Cytokeratin 7 antibody clone KRT7/760 as a reliable reagent for epithelial research and cancer pathology.

The antibody shows distinct cytoplasmic staining in lung, breast, ovary, and urinary tract epithelium. In pathology, it is widely applied to distinguish primary from metastatic tumors, particularly when paired with keratin 20. The CK7/CK20 profile remains a cornerstone of immunohistochemistry in tumor classification.

In oncology, Cytokeratin 7 Antibody KRT7/760 has been applied to research on breast and ovarian cancers. Its detection supports studies into tumor histogenesis, progression, and therapeutic stratification. It has also been used to identify circulating tumor cells, highlighting its value in studies of metastasis and patient monitoring.

The antibody also supports developmental and cell biology studies, where keratin 7 expression provides a marker for epithelial differentiation and cytoskeletal organization.

Validated in tissue and cell-based applications, the antibody consistently produces reproducible cytoplasmic staining with minimal background. Alternate names include CK7 antibody, keratin 7 antibody, and type II cytokeratin antibody.

This mAb is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-).

For an alternative antibody targeting the same protein, see our Cytokeratin 7 Antibody OV-TL12/30 (V2654), a mouse monoclonal antibody validated for WB, IHC, IF, and flow cytometry.

Application Notes

Optimal dilution of the Cytokeratin 7 Antibody KRT7/760 should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant full-length human KRT7 protein was used as the immunogen for the Cytokeratin 7 Antibody KRT7/760.

Storage

Store the Cytokeratin 7 Antibody KRT7/760 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

Keratin 7 antibody, CK7 antibody, CK-7 antibody, KRT7 antibody, Cytokeratin-7 antibody, Keratin Type II Cytoskeletal 7 antibody, Type II Cytokeratin 7 antibody, Cytokeratin 7/CK7 antibody, K7 antibody

Copyright © NSJ Bioreagents. All rights reserved.