


## Cytokeratin 8 Antibody [clone SPM538] (V9038)


Catalog No.	Formulation	Size
V9038-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V9038-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V9038SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V9038IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	SPM538
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P05787
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Flow Cytometry : 1-2ug/10 <sup>6</sup> cells Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This Cytokeratin 8 antibody is available for research use only.




Immunofluorescent staining of fixed human MCF7 cells with Cytokeratin 8 antibody (clone SPM538, green) and Reddot nuclear stain (red).



IHC: Formalin-fixed, paraffin-embedded human lung carcinoma stained with Cytokeratin 8 antibody (clone SPM538).



Flow cytometry testing of MeOH fixed human HeLa cells with Cytokeratin 8 antibody (clone SPM538); Red=isotype control, Blue= Cytokeratin 8 antibody.



SDS-PAGE analysis of purified, BSA-free Cytokeratin 8 antibody (clone SPM538) as confirmation of integrity and purity.

## Description

Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenocarcinomas and ductal carcinomas. It is absent in squamous cell carcinomas. Hepatocellular carcinomas are defined by the use of antibodies that recognize only cytokeratin 8 and 18. CK8 exists on several types of normal and neoplastic epithelia, including many ductal and glandular epithelia such as colon, stomach, small intestine, trachea, and esophagus as well as in transitional epithelium. Anti-CK8 does not react with skeletal muscle or nerve cells. Epithelioid sarcoma, chordoma, and adamantinoma show strong positivity corresponding to that of simple epithelia (with antibodies against CK8, CK18 and CK19). Reportedly, anti-CK8 is useful for the differentiation of lobular ('ring-like, perinuclear') from ductal ('peripheral-predominant') carcinoma of the breast.

## Application Notes

The optimal dilution of the Cytokeratin 8 antibody for each application should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A cytoskeleton preparation containing keratin 8 was used as the immunogen for this Cytokeratin 8 antibody.

## Storage

Store the Cytokeratin 8 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).