
Cytokeratin 8/18 Antibody Cocktail B22.1 + B23.1 / Mouse Monoclonals [clone B22.1 + B23.1] (V3084)

Catalog No.	Formulation	Size
V3084-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3084-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3084SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3084IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Citations (10)

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	B22.1 + B23.1
Purity	Protein G affinity chromatography
UniProt	P05787, P05783
Localization	Cytoplasmic
Applications	1-2ug/10^6 cells1-2ug/ml1-2ug/ml1-2ug/ml for 30 min at RT (1) (2)
Limitations	This Cytokeratin 8/18 Antibody Cocktail B22.1 + B23.1 is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human lung carcinoma stained with Cytokeratin 8/18 Antibody Cocktail B22.1 + B23.1.

Description

Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This mAb cocktail recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the normal epidermis so only Paget cells are stained.

Researchers studying the K8/K18 intermediate filament system can explore our Cytokeratin 8/18 Antibody page for additional reagents targeting these widely expressed epithelial keratins.

Application Notes

Optimal dilution of the Cytokeratin 8/18 Antibody Cocktail B22.1 + B23.1 should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

2. Immunohistochemical staining with this mAb is indistinguishable from that obtained with monoclonal antibody 5D3.

Immunogen

A cytoskeletal preparation from HeLa epitheliod carcinoma cells (B22.1) and PMC-42 breast carcinoma cells of human origin (B23.1) were used as the immunogen for the Cytokeratin 8/18 antibody cocktail.

Storage

Store the Cytokeratin 8/18 antibody cocktail at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.