


## Daxx Antibody [clone PCRP-DAXX-8B7] (V9187)

Catalog No.	Formulation	Size
V9187-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9187-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9187SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	PCRP-DAXX-8B7
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q9UER7
<b>Localization</b>	Nucleus, Cytoplasm
<b>Applications</b>	Flow Cytometry : 1-2ug/million cells Immunofluorescence : 1-2ug/ml
<b>Limitations</b>	This Daxx antibody is available for research use only.


Human Protein Microarray Specificity Validation




Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Daxx antibody (clone PCRP-DAXX-8B7). These results demonstrate the foremost specificity of the PCRP-DAXX-8B7 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD#39;s) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD#39;s) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



Immunofluorescent staining of human U87 cells using Daxx antibody (green, clone PCR-P-DAXX-8B7) and phalloidin (red).



FACS staining of PFA-fixed human HeLa cells with Daxx antibody (blue, clone PCR-P-DAXX-8B7), and unstained cells (red).



SDS-PAGE analysis of purified, BSA-free Daxx antibody (clone PCR-P-DAXX-8B7) as confirmation of integrity and purity.

## Description

Daxx interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, it functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli.

## Application Notes

Optimal dilution of the Daxx antibody should be determined by the researcher.

## Immunogen

Recombinant full-length human DAXX protein was used as the immunogen for the Daxx antibody.

## Storage

Aliquot the Daxx antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

