


# DAZL Antibody [clone DAZL/4253] (V8666)


| Catalog No.    | Formulation  | Size   |
|----------------|--|--------|
| V8666-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8666-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug  |
| V8666SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                             | 100 ug |


# **Bulk quote request**

| Availability       | 1-3 business days   |
|--------------------|---|
| Species Reactivity | Human   |
| Format             | Purified  |
| Clonality          | Monoclonal (mouse origin)   |
| Isotype            | Mouse IgG1  |
| Clone Name         | DAZL/4253   |
| Purity             | Protein G affinity chromatography   |
| UniProt            | Q92904  |
| Localization       | Cytoplasmic, nuclear  |
| Applications       | Immunofluorescence : 1-2ug/ml Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT |
| Limitations        | This DAZL antibody is available for research use only.  |



IHC staining of FFPE human testis with DAZL antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.





SDS-PAGE analysis of purified, BSA-free DAZL antibody as confirmation of integrity and purity.

## **Description**

Specific for human DAZL, a member of the DAZ (deleted in azoospermia) family of RNA binding proteins. DAZL is expressed in fetal and adult testes and ovaries, and is believed to play a role in germ cell development. In adult germ cells, the expression of DAZL is localized to the cytoplasm. Mutations in this gene have been linked to severe spermatogenic failure and infertility in males. Clone DAZL/4253 recognizes DAZL at its C terminus, and is used in detection of DAZL expression by various methods.

### **Application Notes**

Optimal dilution of the DAZL antibody should be determined by the researcher.

#### **Immunogen**

Amino acids CRVHHFRRSRAMLKSV from the human protein (C-terminal region) was used as the immunogen for the DAZL antibody.

#### **Storage**

Store the DAZL antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).