


DCC Antibody (RQ5519)


Catalog No.	Formulation	Size
RQ5519	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P43146
Applications	Western Blot: 0.25-0.5ug/ml Immunohistochemistry (FFPE): 1-2ug/ml Direct ELISA: 0.1-0.5ug/ml
Limitations	This DCC antibody is available for research use only.




Western blot testing of human 1) U-87 MG, 2) U-2 OS, 3) A431 and 4) A549 cell lysate with DCC antibody. Expected molecular weight: 155~190 kDa depending on glycosylation level.




IHC staining of FFPE human glioma with DCC antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE human glioma with DCC antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE mouse brain with DCC antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE rat brain with DCC antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.

Description

Deleted in Colorectal Carcinoma, also known as DCC, is a protein which in humans is encoded by the DCC gene. It is mapped to 18q21.2. This gene encodes a netrin 1 receptor. The transmembrane protein is a member of the immunoglobulin superfamily of cell adhesion molecules, and mediates axon guidance of neuronal growth cones towards sources of netrin 1 ligand. The cytoplasmic tail interacts with the tyrosine kinases Src and focal adhesion kinase (FAK, also known as PTK2) to mediate axon attraction. The protein partially localizes to lipid rafts, and induces apoptosis in the absence of ligand. The protein functions as a tumor suppressor, and is frequently mutated or downregulated in colorectal cancer and esophageal carcinoma.

Application Notes

Optimal dilution of the DCC antibody should be determined by the researcher.

Immunogen

A human recombinant protein (amino acids Q851-F1447) was used as the immunogen for the DCC antibody.

Storage

After reconstitution, the DCC antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.