


DCX Antibody (F48191)

| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F48191-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F48191-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)

| | |
|-----------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Predicted Reactivity | Mouse, Rat |
| Format | Antigen affinity purified |
| Host | Rabbit |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity |
| UniProt | O43602 |
| Localization | Cytoplasmic |
| Applications | Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50 |
| Limitations | This DCX antibody is available for research use only. |



Western blot analysis of human brain tissue lysate using DCX antibody at 1:1000.
Predicted molecular weight: 40-50 kDa.



IHC analysis of FFPE human skeletal muscle tissue stained with DCX antibody

Description

In the developing cortex, cortical neurons must migrate over long distances to reach the site of their final differentiation. DCX is a cytoplasmic protein which appears to direct neuronal migration by regulating the organization and stability of microtubules. It contains two doublecortin domains, which bind microtubules. In addition, this protein interacts with LIS1, the regulatory gamma subunit of platelet activating factor acetylhydrolase, and this interaction is important to proper microtubule function in the developing cortex. Mutations in the gene encoding DCX are a cause of X-linked lissencephaly.

Application Notes

Titration of the DCX antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 107-137 from the human protein was used as the immunogen for this DCX antibody.

Storage

Aliquot the DCX antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.