


# Decorin Antibody / DCN [clone DCN/8760R] (V4308)

Catalog No.	Formulation	Size
V4308-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4308-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4308SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


# Recombinant RABBIT MONOCLONAL

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	DCN/8760R
Purity	Protein A/G affinity
UniProt	P07585
Localization	Secreted
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT
Limitations	This Decorin antibody is available for research use only.



IHC staining of FFPE human prostate carcinoma tissue with Decorin antibody (clone DCN/8760R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



## **Description**

Decorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGFbeta and is a direct or indirect negative modulator of TGFbeta synthesis. Inhibition of Decorin core protein gene expression by the combination of IFN-gamma and TNFAlpha may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.

#### **Application Notes**

Optimal dilution of the Decorin antibody should be determined by the researcher.

#### **Immunogen**

A recombinant human Decorin protein fragment (within amino acids 212-336) was used as the immunogen for the Decorin antibody.

#### **Storage**

Aliquot the Decorin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.