


Decorin Antibody Rabbit Monoclonal / DCN [clone DCN/8760R] (V4308)

Catalog No.	Formulation	Size
V4308-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4308-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4308SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	DCN/8760R
Purity	Protein A/G affinity
UniProt	P07585
Localization	Secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This Decorin antibody rabbit monoclonal is available for research use only.



Immunohistochemistry of Decorin antibody in human prostate carcinoma tissue. Formalin-fixed, paraffin-embedded human prostate carcinoma was stained using rabbit monoclonal Decorin antibody (clone DCN/8760R). Heat induced epitope retrieval was performed by boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 20 min followed by cooling prior to staining. HRP-DAB brown chromogenic signal highlights extracellular stromal connective tissue and desmoplastic matrix surrounding malignant glands, consistent with Decorin localization within collagen-rich tumor stroma.



SDS-PAGE analysis of purified, BSA-free Decorin antibody rabbit monoclonal (clone DCN/8760R) as confirmation of integrity and purity.

Description

Decorin Antibody Rabbit Monoclonal detects Decorin, a small leucine-rich extracellular matrix proteoglycan encoded by the DCN gene and widely expressed in connective tissues. Clone DCN/8760R is a rabbit monoclonal antibody developed for research applications focused on stromal biology, collagen organization, and extracellular matrix remodeling in normal and diseased tissues.

Decorin antibody, also referred to as DCN antibody and small leucine-rich proteoglycan decorin antibody in the literature, recognizes a secreted matrix proteoglycan composed of tandem leucine-rich repeat domains and a single glycosaminoglycan side chain. Decorin binds to fibrillar collagens, particularly type I collagen, where it regulates collagen fibrillogenesis, fiber diameter, and interfibrillar spacing. Through these structural interactions, Decorin contributes to tissue tensile strength and extracellular matrix stability.

In addition to its structural role, Decorin modulates signaling pathways by interacting with growth factors and receptor tyrosine kinases, including transforming growth factor beta and epidermal growth factor receptor. These interactions influence cellular proliferation, migration, and differentiation, positioning Decorin as both a matrix scaffold and a regulatory molecule within the tissue microenvironment. DCN expression is prominent in skin, tendon, ligament, cornea, prostate stroma, placenta, and other collagen-rich tissues. In tissue-based analyses, Decorin typically demonstrates extracellular stromal staining localized between collagen bundles and surrounding glandular structures.

Altered Decorin expression has been implicated in fibrosis, abnormal wound repair, and tumor progression, where extracellular matrix composition shapes stromal signaling and structural organization. Changes in Decorin distribution may accompany desmoplastic reactions and remodeling of collagen architecture in malignancy. A Decorin Antibody Rabbit Monoclonal such as clone DCN/8760R supports investigations into extracellular matrix biology, connective tissue development, fibrosis research, and stromal-tumor interactions. This antibody targets Decorin in research applications and is available from NSJ Bioreagents.

Application Notes

1. Optimal dilution of the Decorin antibody rabbit monoclonal should be determined by the researcher.
2. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.

Immunogen

A recombinant human Decorin protein fragment (within amino acids 212-336) was used as the immunogen for the Decorin antibody rabbit monoclonal.

Storage

Aliquot the Decorin antibody rabbit monoclonal and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

