


Desmoglein-3 Antibody Recombinant Rabbit MAb / DSG3 [clone DSG3/8252R] (V4573)

Catalog No.	Formulation	Size
V4573-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4573-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4573SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	DSG3/8252R
Purity	Protein A/G affinity
UniProt	P32926
Localization	Cell surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Desmoglein-3 antibody is available for research use only.



Immunohistochemistry of Desmoglein-3 antibody in human skin tissue. Formalin-fixed, paraffin-embedded human skin demonstrates membranous staining predominantly in basal and suprabasal keratinocytes, consistent with Desmoglein-3 localization at desmosomal cell-cell junctions within stratified squamous epithelium. Heat-induced epitope retrieval was performed by boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 minutes followed by cooling prior to staining. The recombinant rabbit monoclonal antibody clone DSG3/8252R was used as the primary antibody. Inset: PBS was used in place of the primary antibody as a secondary antibody negative control.



SDS-PAGE analysis of purified, BSA-free DSG3 antibody recombinant rabbit mAb clone DSG3/8252R as confirmation of integrity and purity.

Description

Desmoglein-3 antibody recombinant rabbit mAb clone DSG3/8252R recognizes Desmoglein-3, a calcium-dependent desmosomal cadherin encoded by the DSG3 gene on chromosome 18q12.1. Desmoglein-3, also referred to as DSG3 in the literature, is a single-pass transmembrane glycoprotein localized to desmosomes at the plasma membrane of stratified epithelial cells, where it mediates intercellular adhesion and supports epithelial structural integrity. Desmoglein-3 antibody, also known as DSG3 antibody, is widely used in research focused on epithelial biology, mucosal differentiation, and autoimmune blistering diseases. This recombinant rabbit monoclonal antibody supports detection of membranous DSG3 expression in epithelial tissues.

Desmoglein-3 is a member of the cadherin superfamily and functions as a core component of desmosomal junctions. Its extracellular cadherin repeats mediate calcium-dependent homophilic adhesion between adjacent epithelial cells, while its cytoplasmic domain associates with desmosomal plaque proteins including Plakoglobin and Desmoplakin. Through these interactions, Desmoglein-3 links keratin intermediate filaments to the plasma membrane, reinforcing mechanical stability in tissues exposed to shear and friction, particularly mucosal epithelia and basal epidermal layers.

Expression of DSG3 is predominantly observed in the basal and suprabasal layers of stratified squamous epithelia, including oral mucosa, esophagus, and skin. In contrast to Desmoglein-1, which is more abundant in superficial epidermal layers, Desmoglein-3 is enriched in deeper epithelial layers and plays a critical role in maintaining mucocutaneous adhesion. Autoantibodies directed against Desmoglein-3 are strongly associated with pemphigus vulgaris, where disruption of desmosomal adhesion results in intraepidermal blister formation. Altered DSG3 expression has also been reported in certain squamous cell carcinomas and epithelial malignancies.

Structurally, Desmoglein-3 contains multiple extracellular cadherin domains, a transmembrane region, and an intracellular tail that anchors to the desmosomal plaque complex. Beyond structural adhesion, DSG3 participates in signaling pathways that influence keratinocyte proliferation, differentiation, and epithelial homeostasis. Dysregulated expression may contribute to altered epithelial architecture and tumor progression in stratified epithelia. Through its essential role in desmosomal cohesion, Desmoglein-3 remains central to research in epithelial adhesion biology and mucocutaneous disease mechanisms.

Explore our [Desmoglein 3 Antibody - Human Protein Microarray Validated Clone DSG3/2838](#) page for a broader view of DSG3 expression in stratified epithelia with supporting microarray specificity validation data.

Application Notes

Optimal dilution of the Desmoglein-3 antibody recombinant rabbit mAb should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 300-500) from the human protein was used as the immunogen for the Desmoglein-3 antibody recombinant rabbit mAb.

Storage

Aliquot the Desmoglein-3 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

