
DLL4 Antibody (C-Terminal Region) (F54132)

Catalog No.	Formulation	Size
F54132-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F54132-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	Q9NR61
Applications	WB: 1:1000-1:2000
Limitations	This DLL4 antibody is available for research use only.



Western blot testing of DLL4 antibody at 1:1000: Lane 1) mouse kidney, 2) mouse lung and 3) rat lung lysate. Predicted molecular weight ~75 kDa.

Description

Delta-like protein 4 is involved in the Notch signaling pathway as Notch ligand. Activates NOTCH1 and NOTCH4. Involved in angiogenesis; negatively regulates endothelial cell proliferation and migration and angiogenic sprouting. Essential for retinal progenitor proliferation is required for suppressing rod fates in late retinal progenitors as well as for proper generation of other retinal cell types. During spinal cord neurogenesis, inhibits V2a interneuron fate. [UniProt]

Application Notes

The stated application concentrations are suggested starting points. Titration of the DLL4 antibody may be required due

to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 608-641 from human DLL4 was used as the immunogen for the DLL4 antibody.

Storage

Aliquot the DLL4 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Copyright © NSJ Bioreagents. All rights reserved.