


# **DMRT1 Antibody (R32367)**


Catalog No.	Formulation	Size
R32367	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q9Y5R6
Localization	Nuclear
Applications	Western Blot: 0.1-0.5ug/ml IHC (FFPE): 0.5-1ug/ml
Limitations	This DMRT1 antibody is available for research use only.




Western blot testing of 1) rat testis and 2) mouse testis lysate with DMRT1 antibody. Expected/observed molecular weight ~39 kDa.



IHC testing of FFPE human testis with DMRT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE mouse testis with DMRT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat testis with DMRT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

## **Description**

Doublesex and mab-3 related transcription factor 1, also known as DMRT1, is a protein which in humans is encoded by the DMRT1 gene. This gene is found in a cluster with two other members of the gene family, having in common a zinc finger-like DNA-binding motif (DM domain). The DM domain is an ancient, conserved component of the vertebrate sex-determining pathway that is also a key regulator of male development in flies and nematodes. This gene exhibits a gonad-specific and sexually dimorphic expression pattern. Defective testicular development and XY feminization occur when this gene is hemizygous.

### **Application Notes**

Optimal dilution of the DMRT1 antibody should be determined by the researcher.

#### **Immunogen**

Amino acids RDCQCKKCNLIAERQRVMAAQVALRRQQAQE of human DMRT1 were used as the immunogen for the DMRT1 antibody.

#### **Storage**

After reconstitution, the DMRT1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.