
# Dnmt3b Antibody (RQ6614)


| Catalog No. | Formulation | Size |
|---|---|---|
| RQ6614 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human, Monkey |
| **Format** | Antigen affinity purified |
| **Host** | Rabbit |
| **Clonality** | Polyclonal (rabbit origin) |
| **Isotype** | Rabbit IgG |
| **Purity** | Antigen affinity purified |
| **Buffer** | Lyophilized from 1X PBS with 2% Trehalose |
| **UniProt** | Q9UBC3 |
| **Applications** | Western blot: 1-2ug/ml<br>Flow cytometry: 1-3ug/million cells<br>Direct ELISA: 0.1-0.5ug/ml |
| **Limitations** | This Dnmt3b antibody is available for research use only. |




Flow cytometry testing of human SiHa cells with Dnmt3b antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Dnmt3b antibody.




Western blot testing of 1) human HeLa, 2) monkey COS-7 and 3) human K562 cell lysate with Dnmt3b antibody. Predicted molecular weight: 95 kDa.

## Description

DNA (cytosine-5-)-methyltransferase 3 beta, is an enzyme that in humans in encoded by the DNMT3B gene. CpG methylation is an epigenetic modification that is important for embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a DNA methyltransferase which is thought to function in de novo methylation, rather than maintenance methylation. The protein localizes primarily to the nucleus and its expression is developmentally regulated. Mutations in this gene cause the immunodeficiency-centromeric instability-facial anomalies (ICF) syndrome. Eight alternatively spliced transcript variants have been described. The full length sequences of variants 4 and 5 have not been determined.

## Application Notes

Optimal dilution of the Dnmt3b antibody should be determined by the researcher.

## Immunogen

Recombinant human protein (amino acids K232-K268) was used as the immunogen for the Dnmt3b antibody.

## Storage

After reconstitution, the Dnmt3b antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.


Copyright © NSJ Bioreagents. All rights reserved.