


DR5 Antibody / TRAIL R2 / TNFRSF10B (RQ6120)

Catalog No.	Formulation	Size
RQ6120	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	O14763
Applications	Western Blot : 1-2ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This DR5 antibody is available for research use only.



Western blot testing of human 1) K562, 2) HT1080 and 3) HepG2 lysate with DR5 antibody. Expected molecular weight: ~40 kDa (mature form) and ~48 kDa (precursor). This protein may also be visualized at ~60 kDa.

Description

TNFRSF10B (Tumor necrosis factor receptor superfamily, member 10b) is a human gene. It is also known as DR5, CD262, KILLER, TRICK2, TRICKB, ZTNFR9, TRAILR2, TRICK2A, TRICK2B, TRAIL-R2, KILLER/DR5. The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces apoptosis signal. Mice have a homologous gene, *tnfrsf10b* that has been essential in the elucidation of the function of this gene in humans. Studies with FADD-deficient mice suggested that FADD, a death domain containing

adaptor protein, is required for the apoptosis mediated by this protein. By analysis of radiation hybrid panels, this gene is mapped to chromosome 8p22-p21. Northern blot analysis indicated that TRAILR2 was expressed as a 4.4-kb mRNA in all tissues tested, with the highest levels of expression in peripheral blood lymphocytes, spleen, and ovary.

Application Notes

Optimal dilution of the DR5 antibody should be determined by the researcher.

Immunogen

A human recombinant partial protein (amino acids Q70-K427) was used as the immunogen for the DR5 antibody.

Storage

After reconstitution, the DR5 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.