


dsDNA Antibody [clone DRNC1-1] (V7189)


| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V7189-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V7189-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V7189SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |
| V7189IHC-7ML | Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only* | 7 ml |

[Bulk quote request](#)

| | |
|--------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG3, kappa |
| Clone Name | DRNC1-1 |
| Purity | Protein G affinity chromatography |
| UniProt | Not Known |
| Localization | Nuclear |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT (1) |
| Limitations | This dsDNA antibody is available for research use only. |



IHC testing of FFPE human tonsil with dsDNA antibody (clone DRNC1-1). FFPE testing requires sections to be boiled in pH 9 10mM Tris with 1mM EDTA for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.



IHC testing of FFPE human colon carcinoma with dsDNA antibody (clone DRNC1-1). FFPE testing requires sections to be boiled in pH 9 10mM Tris with 1mM EDTA for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.

Description

Deoxyribonucleic acid (DNA) is a nucleic acid that stores long-term information regarding the development and function of all known living organisms. DNA consists of two long nucleotide polymers, which are composed of four bases, namely adenine, thymine, guanine and cytosine, all of which are flanked by a phosphate-deoxyribose backbone. This mAb can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells.

Application Notes

Titering of the dsDNA antibody may be required for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Cell nuclei were used as the immunogen for the dsDNA antibody.

Storage

Store the dsDNA antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).