


## Dynamin 2 Antibody / DNM2 (RQ4377)

Catalog No.	Formulation	Size
RQ4377	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Mouse, Rat
<b>Predicted Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
<b>UniProt</b>	P50570
<b>Localization</b>	Cytoplasm, cytoskeleton, cell junction
<b>Applications</b>	Western Blot : 0.5-1ug/ml Direct ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This Dynamin 2 antibody is available for research use only.



Western blot testing of 1) rat brain and 2) mouse brain lysate with Dynamin 2 antibody at 0.5ug/ml. Predicted molecular weight ~98 kDa.

## Description

Dynamin-2 is a protein that in humans is encoded by the DNM2 gene. Dynamins represent one of the subfamilies of GTP-binding proteins. These proteins share considerable sequence similarity over the N-terminal portion of the molecule,

which contains the GTPase domain. Dynamins are associated with microtubules. They have been implicated in cell processes such as endocytosis and cell motility, and in alterations of the membrane that accompany certain activities such as bone resorption by osteoclasts. Dynamins bind many proteins that bind actin and other cytoskeletal proteins. Dynamins can also self-assemble, a process that stimulates GTPase activity. Five alternatively spliced transcripts encoding different proteins have been described. Additional alternatively spliced transcripts may exist, but their full-length nature has not been determined.

## Application Notes

Optimal dilution of the Dynamin 2 antibody should be determined by the researcher.

## Immunogen

A recombinant human protein corresponding to amino acids A177-Q441 was used as the immunogen for the Dynamin 2 antibody.

## Storage

After reconstitution, the Dynamin 2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.