


E-Cadherin Antibody / CDH1 [clone SPM381] (V3321)


Catalog No.	Formulation	Size
V3321-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3321-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3321SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM381
Purity	Protein G affinity chromatography
UniProt	P12830
Localization	Cytoplasmic, membranous
Applications	Immunofluorescence : 2-4ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This E-Cadherin antibody is available for research use only.



IHC testing of FFPE human skin with E-Cadherin antibody (clone SPM381). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



Immunofluorescent staining of human MCF7 cells with E-Cadherin antibody (clone SPM381, green) and Reddot nuclear stain (red).

Description

Recognizes a protein of 80-120kDa, identified as E-cadherin/CDH1. Cadherins comprise a family of Ca^{2+} -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH2 terminal repeats. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as β -catenin, to regulate cadherin function. E-cadherin plays an important role in epithelial cell adhesion. A decreased expression of E-cadherin is associated with metastatic potential and poor prognosis in breast cancer, prostate and esophageal cancer. In combination with p120 Catenin, it is useful for the differentiation between ductal (E-cadherin +) and lobular (E-cadherin -) breast carcinomas. It may also help in diagnosis of mesothelioma.

Application Notes

Optimal dilution of the E-Cadherin antibody should be determined by the researcher.

Immunogen

A recombinant human protein was used as the immunogen for the E-Cadherin antibody.

Storage

Store the E-Cadherin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).