



E2F1 Antibody (F50542)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F50542-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F50542-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

Bulk quote request

| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity |
| UniProt | Q01094 |
| Applications | Western Blot: 1:1000 |
| Limitations | This E2F1 antibody is available for research use only. |







Description

E2F1 is a transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F1 binds preferentially RB1 in a cell-cycle dependent manner. It can mediate both cell proliferation and TP53/p53-dependent apoptosis. Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters. [UniProt]

Application Notes

Titration of the E2F1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 342-371 from the human protein was used as the immunogen for this E2F1 antibody.

Storage

Aliquot the E2F1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.