


EAAT1 Antibody / Excitatory amino acid transporter 1 / GLAST (RQ7320)


Catalog No.	Formulation	Size
RQ7320	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P43003
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml
Limitations	This EAAT1 antibody is available for research use only.




Immunofluorescent staining of FFPE rat brain tissue with EAAT1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.



Immunofluorescent staining of FFPE mouse brain tissue with EAAT1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.




IHC staining of FFPE rat cerebellum tissue with EAAT1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat brain tissue with EAAT1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat brain tissue with EAAT1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE mouse brain tissue with EAAT1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) rat brain and 2) mouse brain tissue lysate with EAAT1 antibody. Predicted molecular weight ~60 kDa.

Description

Excitatory amino acid transporter 1 (EAAT1), also called GLAST, is a protein that, in humans, is encoded by the SLC1A3 gene. It is mapped to 5p13.2. This gene encodes a member of a high affinity glutamate transporter family. This gene functions in the termination of excitatory neurotransmission in central nervous system. Mutations are associated with episodic ataxia, Type 6. Alternative splicing results in multiple transcript variants.

Application Notes

Optimal dilution of the EAAT1 antibody should be determined by the researcher.

Immunogen

Amino acids EMKKPYQLIAQDNETEKPID were used as the immunogen for the EAAT1 antibody.

Storage

After reconstitution, the EAAT1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.