


# EBV Antibody Cocktail / Epstein-Barr Virus / LMP-1 [clone CS1-4] (V8335)

Catalog No.	Formulation	Size
V8335-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8335-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8335SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	EBV
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CS1-4
Purity	Protein G affinity chromatography
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This EBV antibody cocktail is available for research use only.



IHC staining of FFPE Hodgkin's Lymphoma with EBV antibody cocktail (clone CS1-4). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.

## **Description**

This antibody cocktail is a mixture of four different monoclonal antibodies (CS1 + CS2 + CS3 + CS4). It is specific to 60kDa latent membrane protein (LMP-1) encoded by the BNLF1 gene of the EBV. Each clone reacts with different epitopes on the hydrophilic C-terminus of the cytoplasmic domain of LMP-1. The cocktail stains strongly with EBV-positive lymphoblastoid cell lines and EBV infected B cell immunoblasts in infectious mononucleosis. EBV, also designated

human herpesvirus 4 (HHV-4), is a member of the herpesvirus family and is one of the most common human viruses. EBV infects B cells and, though often asymptomatic, it can cause infectious mononucleosis, a disease characterized by fatigue, fever, sore throat and muscle soreness.

#### **Application Notes**

Optimal dilution of the EBV antibody cocktail should be determined by the researcher.

## **Immunogen**

Recombinant fusion protein containing the sequence of bacterial beta-galactosidase and the carboxyl half of EBV-encoded LMP was used as the immunogen for the EBV antibody cocktail.

# **Storage**

Store the EBV antibody cocktail at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).