


# EG-VEGF Antibody / Prokineticin-1 (R32153)

Catalog No.	Formulation	Size
R32153	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P58294
Applications	Western Blot : 0.1-0.5ug/ml ELISA : 0.1-0.5ug/ml (human protein tested); request BSA-free format for coating
Limitations	This EG-VEGF antibody is available for research use only.



# **Description**

Prokineticin-1 is also known as EG-VEGF. The International Radiation Hybrid Mapping Consortium mapped the PROK1 gene to chromosome 1. The protein encoded by this gene induces proliferation, migration, and fenestration (the formation of membrane discontinuities) in capillary endothelial cells derived from endocrine glands. It has little or no effect on a variety of other endothelial and non-endothelial cell types. Its expression is restricted to the steroidogenic glands (ovary, testis, adrenal, and placenta), is induced by hypoxia, and often complementary to the expression of vascular endothelial growth factor (VEGF), suggesting that these molecules function in a coordinated manner.

# **Application Notes**

Optimal dilution of the EG-VEGF antibody should be determined by the researcher.

#### **Immunogen**

Amino acids 20-105 of human EG-VEGF were used as the immunogen for the EG-VEGF antibody.

# **Storage**

After reconstitution, the EG-VEGF antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.