


# ELK1 Antibody / ETS Like-1 protein Elk-1 [clone PCRP-ELK1-1B9] (V9226)


Catalog No.	Formulation	Size
V9226-100UG	0.2~mg/ml in 1X PBS with $0.1~mg/ml$ BSA (US sourced), $0.05%$ sodium azide	100 ug
V9226-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9226SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a
Clone Name	PCRP-ELK1-1B9
Purity	Protein A/G affinity
UniProt	P19419
Localization	Nucleus
Applications	Immunoprecipitation: 1-2ug per 100-500ug of total protein (1ml of cell lysate)
Limitations	This ELK1 antibody is available for research use only.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using ELK1 antibody (clone PCRP-ELK1-1B9). These results demonstrate the foremost specificity of the PCRP-ELK1-1B9 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free ELK1 antibody (clone PCRP-ELK1-1B9) as confirmation of integrity and purity.

# **Description**

Ets-1 is the prototype member of a family of genes identified on the basis of homology to the v-Ets oncogene isolated from the E26 erythroblastosis virus. This family of genes currently includes Ets-1, Ets-2, Erg-1 3, Elk-1, Elf-1, Elf-5, NERF, PU.1, PEA3, ERM, FEV, ER8I, Fli-1, TEL, Spi-B, ESE-1, ESE-3A, Net, ABT1and ERF. Members of the Ets gene family exhibit varied patterns of tissue expression, and share a highly conserved carboxy-terminal domain containing a sequence related to the SV40 large T antigen nuclear localization signal sequence. This conserved domain is essential for Ets-1 binding to DNA and is likely to be responsible for the DNA binding activity of all members of the Ets gene family. Several of these proteins have been shown to recognize similar motifs in DNA that share a centrally located 5 element.

### **Application Notes**

Optimal dilution of the ELK1 antibody should be determined by the researcher.

### **Immunogen**

Recombinant full-length human ELK1 protein was used as the immunogen for the ELK1 antibody.

# **Storage**

Aliquot the ELK1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.