

EME1 Antibody [clone MTA31 7H2/1] (V8532)

Catalog No.	Formulation	Size
V8532-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8532-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8532SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	MTA31 7H2/1
Purity	Protein G affinity chromatography
UniProt	Q96AY2
Localization	Nucleus
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This EME1 antibody is available for research use only.



Description

Essential meiotic endonuclease 1 (Eme1), a member of the Eme1/Mms4 family, associates with Mus81 to constitute a heterodimeric endonuclease that has been implicated in mitotic and meiotic recombination in eukaryotes. The Mus81-Eme1 complex cleaves branched DNA structures, especially those arising during stalled DNA replication such as replication forks and 3 DNA flaps. When purified from yeast, this complex cleaves synthetic Holliday junctions into linear duplex DNA. These findings provide compelling evidence that Mus81-Eme1 complexes are essential elements of the eukaryotic nuclear Holliday junction resolvase. Eme1 may also be required in mitosis for the processing of collapsed replication forks. Eme1 is typically localized to the nucleolus and is recruited to regions of DNA damage in S phase cells.

Application Notes

Optimal dilution of the EME1 antibody should be determined by the researcher.

Immunogen

His-tagged recombinant Eme1 of human origin was used as the immunogen for the EME1 antibody.

Storage

Store the EME1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).