


Endoglin Antibody / CD105 [clone ENG/1621] (V8568)


Catalog No.	Formulation	Size
V8568-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8568-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8568SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	ENG/1621
Purity	Protein G affinity chromatography
UniProt	P17813
Localization	Cell surface
Applications	ELISA : order Ab without BSA for coating Flow Cytometry : 1-2ug/million cells Immunofluorescence : 2-4ug/ml
Limitations	This Endoglin antibody is available for research use only.



Flow cytometry testing of PFA-fixed human HeLa cells with Endoglin antibody (clone ENG/1621); Red=isotype control, Blue= Endoglin antibody.



Immunofluorescent staining of PFA-fixed human HeLa cells with Endoglin antibody (clone ENG/1621).



SDS-PAGE analysis of purified, BSA-free Endoglin antibody (clone ENG/1621) as confirmation of integrity and purity.

Description

CD105/Endoglin is a Type I membrane glycoprotein located on cell surfaces and is part of the TGF-beta receptor complex. This protein has been found on endothelial cells, activated macrophages, fibroblasts, and smooth-muscle cells. Endoglin has a role in the development of the cardiovascular system and in vascular remodeling. Its expression is regulated during heart development. CD105 is highly expressed in endothelial cells during tumor angiogenesis and inflammation, with weak or negative expression in vascular endothelium of normal tissues. Angiogenesis is a promising prognostic marker in a variety of tumors. Endoglin is a more specific and sensitive marker for tumor angiogenesis than CD31 or CD34, as it labels only newly-formed blood vessels and may serve as a prognostic marker for Prostate Adenocarcinoma, and cancers of the lung, stomach, breast, and brain.

Application Notes

Optimal dilution of the Endoglin antibody should be determined by the researcher.

Immunogen

A recombinant extracellular fragment from the human protein was used as the immunogen for the Endoglin antibody.

Storage

Store the Endoglin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).