
# Eotaxin 3 Antibody / CCL26 (C-Terminal Region) (F54151)

| Catalog No. | Formulation | Size |
|---|---|---|
| F54151-0.2ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.2 ml |
| F54151-0.05ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.05 ml |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Antigen affinity purified |
| **Host** | Rabbit |
| **Clonality** | Polyclonal (rabbit origin) |
| **Isotype** | Rabbit Ig |
| **Purity** | Antigen affinity |
| **UniProt** | Q9Y258 |
| **Applications** | |
| **Limitations** | This Eotaxin 3 antibody is available for research use only. |




Western blot testing of human 1) 293T/17, 2) HUVEC, 3) K562, 4) NCI-H460 and 5) PC-3 cell lysate with Eotaxin-3 antibody at 1:2000. Predicted molecular weight ~11 kDa.

## Description

Eotaxin 3 is a chemoattractant for eosinophils and basophils and acts as a ligand for C-C chemokine receptor CCR3 which triggers Ca2+ mobilization in eosinophils. [UniProt]

Explore our Eotaxin-3 Antibody page for additional information on this eosinophil chemoattractant involved in allergic inflammation, immune signaling, and tumor-associated inflammatory responses.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the Eotaxin 3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 58-92 from human Eotaxin 3 was used as the immunogen for the Eotaxin 3 antibody.

## Storage

Aliquot the Eotaxin 3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.


Copyright © NSJ Bioreagents. All rights reserved.