


## Epstein-Barr virus induced 3 Antibody / EB13 [clone EB13/8906] (V4606)


Catalog No.	Formulation	Size
V4606-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4606-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4606SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	EB13/8906
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q14213
<b>Localization</b>	Secreted
<b>Applications</b>	ELISA (Order BSA-free Format For Coating) : Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 2-4ug/ml
<b>Limitations</b>	This Epstein-Barr virus induced 3 antibody is available for research use only.



Western blot testing of human placental tissue lysate with EB13 antibody (clone EB13/8906). Predicted molecular weight ~25 kDa but may be observed at higher molecular weights due to glycosylation.



IHC staining of FFPE human placental tissue with Epstein-Barr virus induced 3 antibody (clone EBI3/8906). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free EBI3 antibody (clone EBI3/8906) as confirmation of integrity and purity.

## Description

Epstein-Barr virus-induced gene 3 (Ebi3) is a widely expressed homolog to the interleukin IL-12 p40 subunit protein that forms a heterodimer with either IL-12 p35 or an IL-12 p35 homolog, p28, to create a new cytokine (IL-27). IL-27 is an early product of activated antigen-presenting cells and drives rapid clonal expansion of naive but not memory CD4<sup>+</sup> T cells. Interferon- $\gamma$  differentially regulates expression of the IL-12 family members p35, p40, p19 and Ebi3 in activated human dendritic cells. Ebi3 may function to antagonize IL-12 and to inhibit the development of a Th1 immune response. Ebi3 is strongly expressed in Hodgkin and Reed-Sternberg cells, independently of the EBV status of the tumor cells. Research suggests that Ebi3 may be an additional component of the repertoire employed by Hodgkin and Reed-Sternberg cells to inhibit an effective anti-tumor or anti-viral immune response. The human Ebi3 gene maps to chromosome 19p13.3 and encodes a secreted glycoprotein that is expressed in spleen and tonsils, and at high levels in full-term placenta.

## Application Notes

Optimal dilution of the Epstein-Barr virus induced 3 antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 366-466) from the human protein was used as the immunogen for the Epstein-Barr virus induced 3 antibody.

## Storage

Aliquot the Epstein-Barr virus induced 3 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.