


## ErbB2 Antibody (F50600)


| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F50600-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F50600-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)


|                             |  |
|-----------------------------|--|
| <b>Availability</b>         | 1-3 business days  |
| <b>Species Reactivity</b>   | Human  |
| <b>Predicted Reactivity</b> | Mouse, Rat   |
| <b>Format</b>               | Purified   |
| <b>Host</b>                 | Rabbit   |
| <b>Clonality</b>            | Polyclonal (rabbit origin)   |
| <b>Isotype</b>              | Rabbit Ig  |
| <b>Purity</b>               | Purified   |
| <b>UniProt</b>              | P04626   |
| <b>Applications</b>         | Western Blot : 1:1000<br>Flow Cytometry : 1:10-1:50<br>IHC (Paraffin) : 1:50-1:100 |
| <b>Limitations</b>          | This ErbB2 antibody is available for research use only.                            |



Western blot analysis of ErbB2 in T47D cell lysate, either noninduced (Lane 1) or induced with HRG (2).



Flow cytometric analysis of MCF-7 cells using HER2 / ErbB2 antibody (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.



IHC analysis of FFPE human breast carcinoma tissue stained with ErbB2 antibody

## Description

ErbB2, a member of the EGF receptor family, is an essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. This protein is not activated by EGF, TGF-alpha and amphiregulin. ErbB2 potentially forms a heterodimer with each of the other ERBB receptors. An interaction with PRKCABP has been suggested. Ligand-binding to this Type I membrane protein may increase phosphorylation on tyrosine residues

## Application Notes

Titration of the ErbB2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 21-52 from the human protein was used as the immunogen for this ErbB2 antibody.

## Storage

Aliquot the ErbB2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.