

## Estrogen Inducible Protein pS2 Antibody / TFF1 [clone SPM313] (V2879)


Catalog No.	Formulation	Size
V2879-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2879-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2879SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2879IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml




Citations (2)

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	SPM313
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P04155
<b>Localization</b>	Cytoplasmic, secreted
<b>Applications</b>	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
<b>Limitations</b>	This Estrogen Inducible Protein pS2 antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human ovarian carcinoma stained with Estrogen Inducible Protein pS2 antibody (SPM313).



SDS-PAGE analysis of purified, BSA-free Estrogen Inducible Protein pS2 antibody (clone SPM313) as confirmation of integrity and purity.

## Description

It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.

## Application Notes

Optimal dilution of the Estrogen Inducible Protein pS2 antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

Amino acids CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the human protein were used as the immunogen for the Estrogen Inducible Protein pS2 antibody.

## Storage

Store the Estrogen Inducible Protein pS2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).