


## FAM125A Antibody (N-Terminal Region) (F54126)

Catalog No.	Formulation	Size
F54126-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F54126-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q96EY5
<b>Applications</b>	Western Blot : 1:2000
<b>Limitations</b>	This FAM125A antibody is available for research use only.



Western blot testing of FAM125A antibody at 1:2000 + human PC-3 cell lysate.  
Predicted molecular weight ~29 kDa.

### Description

Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies. May be involved in the ligand-mediated internalization and down-regulation of EGF receptor. [UniProt]

### Application Notes

The stated application concentrations are suggested starting points. Titration of the FAM125A antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## **Immunogen**

A portion of amino acids 16-50 from human FAM125A was used as the immunogen for the FAM125A antibody.

## **Storage**

Aliquot the FAM125A antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.