


## FCGR1A Antibody / CD64 [clone FCGR1A/7497] (V4724)


Catalog No.	Formulation	Size
V4724-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4724-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4724SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2, kappa
Clone Name	FCGR1A/7497
Purity	Protein A/G affinity
UniProt	P12314
Localization	Cell Membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This FCGR1A antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with FCGR1A antibody (clone FCGR1A/7497) at 2ug/ml. Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free FCGR1A antibody (clone FCGR1A/7497) as confirmation of integrity and purity.

## Description

CD64/FCGR1A is a surface glycoprotein with high affinity for monomeric IgG, is expressed constitutively on monocytes and macrophages, and can be induced in neutrophils subsequent to IFN-gamma stimulation. It plays a putative role in the initiation of cell-mediated cytotoxicity. Thus far, three genes encoding four distinct CD64 transcripts have been described. CD64/FCGR1A has been shown to associate with signal transducing subunit of the high affinity IgE receptor.

## Application Notes

Optimal dilution of the FCGR1A antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 1-200) from the human protein was used as the immunogen for the FCGR1A antibody.

## Storage

Aliquot the FCGR1A antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.