


FGF8 Antibody (R30317)


Catalog No.	Formulation	Size
R30317	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
UniProt	P55075
Localization	Secreted
Applications	Western Blot : 0.5-1ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This FGF8 antibody is available for research use only.



Western blot testing of FGF8 antibody and rat ovary tissue lysate. Predicted molecular weight: $\sim\!25$ kDa.



IHC-P: FGF8 antibody testing of rat ovary tissue. HIER: steamed with pH6 citrate buffer.

Description

Fibroblast growth factor 8 (androgen-induced), also known as FGF8 or AIGF, is a human gene which maps to 10q24. The protein encoded by this gene are secreted proteins that interact with FGF tyrosine kinase receptors to mediate growth and development. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogensis. The temporal and spatial patterns of this gene expression suggest that FGF8 is involved in gastrulation, regionalization of the brain, and organogenesis of the limb and face as an embryonic epithelial factor. The adult expression of FGF8 is restricted to gonads, including testes and ovaries. FGF8 stimulated growth of human prostate carcinoma cells and mouse fibroblasts and mammary carcinoma cells in a dose-dependent manner. It also may play an important role in growth and patterning of limbs, face, and central nervous system. FGF8 is expressed in increased levels in breast cancer and in lactating human breast; it was also detected in human milk. A survey of other normal tissues showed that FGF8 is expressed in the proliferative cells of the skin and epithelial cells in colon, ovary, fallopian tube, and uterus.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the FGF8 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the C-terminus of human FGF8 (FMKRLPRGHHTTEQSLRFEFLNY) was used as the immunogen for this FGF8 antibody (100% homologous in human, mouse and rat).

Storage

After reconstitution, the FGF8 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.