


Fibronectin Antibody [clone SPM539] (V9020)

Catalog No.	Formulation	Size
V9020-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V9020-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V9020SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM539
Purity	Protein G affinity chromatography
UniProt	P02751
Localization	Connective tissue matrix
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Fibronectin antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human pancreatic adenocarcinoma stained with Fibronectin antibody (SPM539).

Description

Fibronectin is a dimeric glycoprotein of 440kDa which is present in cells, extracellular matrix, and blood. It possesses at least four binding sites for collagen, glycosaminoglycans, transglutaminase, and a cell surface receptor. Fibronectin is involved in cell adhesion, tissue organization, and wound healing. This mAb is directed against the peptide core and reacts with both the plasma and cellular forms of fibronectin.

Application Notes

The optimal dilution of the Fibronectin antibody for each application should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9 for 10-20 min followed by cooling at RT for 20 minutes.

Immunogen

Purified human fibronectin was used as the immunogen for this Fibronectin antibody.

Storage

Store the Fibronectin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).