


Filaggrin Antibody [clone SPM181] (V3364)

Catalog No.	Formulation	Size
V3364-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3364-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3364SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


 [Citations \(15\)](#)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM181
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
UniProt	P20930
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Filaggrin antibody is available for research use only.



IHC testing of FFPE human skin with Filaggrin antibody (clone SPM181). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



SDS-PAGE analysis of purified, BSA-free Filaggrin antibody (clone SPM181) as confirmation of integrity and purity.

Description

Filaggrin (filament aggregating protein) is a filament-associated protein that binds to keratin fibers in epithelial cells. Ten to twelve filaggrin units are post-translationally hydrolyzed from a large profilaggrin precursor protein during terminal differentiation of epidermal cells. In humans, profilaggrin is encoded by the FLG gene, which is part of the S100 fused-type protein (SFTP) family within the epidermal differentiation complex on chromosome 1q21.

Filaggrin is essential for the regulation of epidermal homeostasis. Within the stratum corneum, filaggrin monomers can become incorporated into the lipid envelope, which is responsible for the skin barrier function. Alternatively, these proteins can interact with keratin intermediate filaments. Filaggrin undergoes further processing in the upper stratum corneum to release free amino acids that assist in water retention. [Wiki]

Application Notes

Optimal dilution of the Filaggrin antibody should be determined by the researcher.

Immunogen

A recombinant human FLG protein was used as the immunogen for this Filaggrin antibody.

Storage

Store the Filaggrin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).