


Flt3 ligand Antibody / Flt3l (R32965)


Catalog No.	Formulation	Size
R32965	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	A0A0G2K6K8
Applications	Western Blot : 0.5-1ug/ml Direct ELISA (recombinant Rat Protein) : 0.1-0.5ug/ml (BSA-free format available)
Limitations	This Flt3 ligand antibody is available for research use only.



Western blot testing of 1) mouse kidney, 2) mouse spleen and 3) rat skeletal muscle tissue lysate with Flt3 ligand antibody at 0.5ug/ml. Predicted molecular weight ~26 kDa.



Western blot testing of 1) rat thymus and 2) mouse thymus tissue lysate with Flt3 ligand antibody at 0.5ug/ml. Predicted molecular weight ~26 kDa.

Description

FLT3LG (FMS-Related Tyrosine Kinase 3 Ligand) also called FLT3 ligand, FL or FLT3L, is a protein which in humans is encoded by the FLT3LG gene. FLT3LG controls the development of DCs and is particularly important for plasmacytoid DCs and CD8-positive classical DCs and their CD103-positive tissue counterparts. Flt3 ligand (FL) is a hematopoietic four helical bundle cytokine. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors. Lyman et al. (1993) found that Flt3 ligand stimulated proliferation of hematopoietic progenitor cells isolated from mouse fetal liver or adult mouse bone marrow. Hannum et al. (1994) concluded that FL enhances the response of stem and primitive progenitor cells to other growth factors to generate all myeloid lineages except erythroid cells. Assays of C-terminally truncated FL proteins confirmed that the N-terminal half conferred FL activity. Depletion of the Pi3k -Mtor negative regulator Pten facilitated Flt3L-driven DC development in culture.

Application Notes

Optimal dilution of the Flt3 ligand antibody should be determined by the researcher.

Immunogen

A recombinant rat protein corresponding to amino acids T28-Q189 was used as the immunogen for the Flt3 ligand antibody.

Storage

After reconstitution, the Flt3 ligand antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.