


FMO2 Antibody / Flavin-containing monooxygenase 2 (R32311)

| Catalog No. | Formulation | Size |
|-------------|---|--------|
| R32311 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

[Bulk quote request](#)

| | |
|---------------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | Human, Mouse, Rat |
| Format | Antigen affinity purified |
| Host | Rabbit |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Antigen affinity |
| Buffer | Lyophilized from 1X PBS with 2% Trehalose |
| UniProt | Q99518 |
| Applications | Western Blot : 0.5-1ug/ml |
| Limitations | This FMO2 antibody is available for research use only. |



Western blot testing of 1) rat lung and 2) mouse lung tissue lysate with FMO2 antibody. Predicted molecular weight ~54 kDa.

Description

Dimethylaniline monooxygenase [N-oxide-forming] 2 is an enzyme that in humans is encoded by the FMO2 gene (flavin containing monooxygenase 2). It is an NADPH-dependent enzyme that catalyzes the N-oxidation of some primary alkylamines through an N-hydroxylamine intermediate. However, some human populations contain an allele (FMO2*2A) with a premature stop codon, resulting in a protein that is C-terminally-truncated, has no catalytic activity, and is likely degraded rapidly. This gene is found in a cluster with other related family members on chromosome 1. Alternative splicing results in multiple transcript variants.

Application Notes

Optimal dilution of the FMO2 antibody should be determined by the researcher.

Immunogen

Amino acids FPNFLHNSKLLEYFRIFAKKFDLLKYIQFQTTVLSVRK of human FMO2 were used as the immunogen for the FMO2 antibody.

Storage

After reconstitution, the FMO2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.