


## Fms-like tyrosine kinase 3 Antibody / FLT3 [clone FLT3/9886] (V5706)


Catalog No.	Formulation	Size
V5706-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5706-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5706SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG
<b>Clone Name</b>	FLT3/9886
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P36888
<b>Localization</b>	Cytoplasm, Membrane
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This Fms-like tyrosine kinase 3 antibody is available for research use only.



IHC staining of FFPE human brain tissue with Fms-like tyrosine kinase 3 antibody (clone FLT3/9886). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Fms-like tyrosine kinase 3 antibody (clone FLT3/9886) as confirmation of integrity and purity.

## Description

Stem cell tyrosine kinase (STK-1) has been cloned from a CD34+ hematopoietic stem cell enriched library and identified as the human homolog of a previously identified gene of Mouse origin designated either Flk-2 or Flt-3. The STK-1 cDNA encodes a protein of 993 amino acids with 85% identity to Flt-3/Flk-2. STK-1 is a member of the type III receptor tyrosine kinase family that includes Kit (steel factor receptor), Fms and PDGF. STK-1 expression in blood and marrow is restricted to CD34+ cells, a population greatly enriched for hematopoietic stem/progenitor cells. STK-1 antiserum recognizes two polypeptides in these cells. The Mouse homolog of STK-1, designated Flt-3/Flk-2, is expressed at high levels in hematopoietic cells and also in neural, gonadal, hepatic and placental tissues. It has been suggested that STK-1 and its murine homolog Flt-3/Flk-2 may function as growth factor receptors on hematopoietic stem and/or progenitor cells.

## Application Notes

Optimal dilution of the Fms-like tyrosine kinase 3 antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 500-800 from human Fms-like tyrosine kinase 3 protein was used as the immunogen for the FLT3 antibody.

## Storage

Aliquot the Fms-like tyrosine kinase 3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.