


Forkhead Box I1 Antibody / FOXI1 [clone PCRP-FOXI1-1C4] (V4449)


Catalog No.	Formulation	Size
V4449-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4449-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4449SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a
Clone Name	PCRP-FOXI1-1C4
Purity	Protein A/G affinity
UniProt	Q12951
Localization	Nucleus
Applications	Flow Cytometry : 1-2ug/million cells Immunofluorescence : 1-2ug/ml Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This Forkhead Box I1 antibody is available for research use only.



Flow cytometry testing of PFA-fixed human HeLa cells with Forkhead Box I1 antibody (clone PCRP-FOXI1-1C4) followed by goat anti-mouse IgG-CF488 (blue), Red = unstained cells.



SDS-PAGE analysis of purified, BSA-free Forkhead Box I1 antibody (clone PCRP-FOXI1-1C4) as confirmation of integrity and purity.



Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using Forkhead Box I1 antibody (clone PCRP-FOXI1-1C4). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.

Description

FOXI1 is a transcription factor required for proper proliferation and differentiation in the gastrointestinal epithelium. [UniProt]

Application Notes

Optimal dilution of the Forkhead Box I1 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 104-258) from the human protein was used as the immunogen for the Forkhead Box I1 antibody.

Storage

Aliquot the Forkhead Box I1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.