


# FOXA2 Antibody (F52908)

Catalog No.	Formulation	Size
F52908-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F52908-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	Q9Y261
Applications	Western Blot : 1:1000
Limitations	This FOXA2 antibody is available for research use only.



Western blot analysis of lysate from (1) HepG2, (2) HT-29, and (3) SW620 cell line using FOXA2 antibody at 1:1000. Predicted molecular weight: 50 kDa.

## **Description**

Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence

5'-[AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). In embryonic development is required for notochord formation. Involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas and lungs; FOXA1 and FOXA2 seem to have at least in part redundant roles. Originally described as a transcription activator for a number of

liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; regulates the expression of genes important for glucose sensing in pancreatic beta-cells and glucose homeostasis. Involved in regulation of fat metabolism. Binds to fibrinogen beta promoter and is involved in IL6-induced fibrinogen beta transcriptional activation. [UniProt]

#### **Application Notes**

Titration of the FOXA2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

This FOXA2 antibody was produced from a rabbit immunized with a KLH conjugated synthetic peptide between 355-389 amino acids from the C-terminal region of human FOXA2.

#### **Storage**

Aliquot the FOXA2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.