


FOXL2 Antibody (RQ6867)


Catalog No.	Formulation	Size
RQ6867	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P58012
Localization	Nuclear
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This FOXL2 antibody is available for research use only.




IHC staining of FFPE mouse ovary tissue with FOXL2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat ovary tissue with FOXL2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




Western blot testing of 1) human Jurkat, 2) rat ovary and 3) mouse ovary tissue lysate with FOXL2 antibody. Predicted molecular weight ~38 kDa, but can be observed at 45-50 kDa.



Flow cytometry testing of fixed and permeabilized human ThP-1 cells with FOXL2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= FOXL2 antibody.



IHC staining of FFPE human ovarian cancer tissue with FOXL2 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lung cancer tissue with FOXL2 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

FOXL2 (Forkhead box protein L2) is a transcription factor belonging to the forkhead/winged-helix family, characterized by a conserved DNA-binding domain that regulates gene expression during development and tissue maintenance. FOXL2 is predominantly expressed in ovarian granulosa cells and certain ocular tissues, where it plays key roles in cell differentiation, growth regulation, and long-term tissue homeostasis.

Functionally, FOXL2 is involved in pathways that influence folliculogenesis, hormone signaling, and extracellular matrix remodeling. It also participates in broader transcriptional networks that coordinate cell identity and developmental timing. Due to its specific expression profile, FOXL2 serves as an important marker in developmental biology and reproductive

research.

The **FOXL2 antibody** enables precise detection of endogenous FOXL2 in applications such as western blot, immunohistochemistry, and immunofluorescence. Researchers use the FOXL2 antibody from NSJ Bioreagents to measure protein levels, determine subcellular localization, and study transcription factor activity under defined experimental conditions. With high specificity and consistent performance, the FOXL2 antibody supports detailed investigations into transcriptional regulation, reproductive biology, and developmental gene networks.

Application Notes

Optimal dilution of the FOXL2 antibody should be determined by the researcher.

Immunogen

Amino acids ACARQPELAMMHCSYWDHD from the human protein were used as the immunogen for the FOXL2 antibody.

Storage

After reconstitution, the FOXL2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.