


# **GABARAP Antibody (F46304)**


| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F46304-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F46304-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

# **Bulk quote request**


| Availability         | 1-3 business days   |
|----------------------|---|
| Species Reactivity   | Human   |
| Predicted Reactivity | Mouse, Rat, Bovine, Rabbit  |
| Format               | Antigen affinity purified   |
| Clonality            | Polyclonal (rabbit origin)  |
| Isotype              | Rabbit Ig   |
| Purity               | Antigen affinity  |
| UniProt              | O95166  |
| Applications         | Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50 Immunofluorescence : 1:10-1:50 |
| Limitations          | This GABARAP antibody is available for research use only.                       |




Western blot testing of GABARAP antibody and Hela cells treated with 26S proteasome complex blocker MG132.




Western blot analysis of GABARAP antibody and 293 cell lysate either nontransfected (Lane 1) or transiently transfected with the GABARAP gene (2).



Fluorescent image of Chloroquine treated U251 cells stained with GABARAP antibody. Alexa Fluor 488 secondary (green) was used. Nuclei were counterstained with Hoechst 33342 (blue) (10 ug/ml, 5 min). GABARAP immunoreactivity is localized to autophagic vacuoles in the cytoplasm.



GABARAP antibody testing of mouse MEFs and mouse motor neurons. Antibodies used were GABARAP antibody (AF488 conjugated) and anti-Tuj1 (AF647 conjugated).



IHC analysis of FFPE human hepatocarcinoma tissue stained with GABARAP antibody

## **Description**

Gamma-aminobutyric acid A receptors [GABA(A) receptors] are ligand-gated chloride channels that mediate inhibitory neurotransmission. GABARAP is GABA(A) receptor-associated protein, which is highly positively charged in its N-terminus and shares sequence similarity with light chain-3 of microtubule-associated proteins 1A and 1B. This protein clusters neurotransmitter receptors by mediating interaction with the cytoskeleton.

## **Application Notes**

Titration of the GABARAP antibody may be required due to differences in protocols and secondary/substrate sensitivity.

### **Immunogen**

A portion of amino acids 1-30 from the human protein was used as the immunogen for this GABARAP antibody.

### **Storage**

Aliquot the GABARAP antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.