


## GAD2 Antibody / GAD65 (F54577)


Catalog No.	Formulation	Size
F54577-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54577-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity purified
<b>UniProt</b>	Q05329
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 1:500-1:2000 Immunohistochemistry (FFPE) : 1:25 Immunofluorescence : 1:25
<b>Limitations</b>	This GAD2 antibody is available for research use only.




Western blot testing of human Jurkat cell lysate with GAD2 antibody. Predicted molecular weight ~65 kDa.



Western blot testing of rat brain tissue lysate with GAD2 antibody. Predicted molecular weight ~65 kDa.



IHC testing of FFPE human pancreas tissue with GAD2 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Immunofluorescent staining of human HEK293 cells with GAD2 antibody (green) and DAPI nuclear stain (blue).

## Description

This gene encodes one of several forms of glutamic acid decarboxylase, identified as a major autoantigen in insulin-dependent diabetes. The enzyme encoded is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantibody and an autoreactive T cell target in insulin-dependent diabetes. This gene may also play a role in the stiff man syndrome. Alternative splicing results in multiple transcript variants that encode the same protein.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the GAD2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 109-138 from the human protein was used as the immunogen for the GAD2 antibody.

## Storage

Aliquot the GAD2 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

