


GAD65 Antibody / GAD2 [clone GAD2/1960] (V3847)

Catalog No.	Formulation	Size
V3847-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3847-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3847SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	GAD2/1960
Purity	Protein G affinity chromatography
UniProt	Q05329
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 0.1-0.2ug/ml for 30 min at RT
Limitations	This GAD65 antibody is available for research use only.




GAD65 Antibody Pancreas IHC. Immunohistochemistry testing of FFPE human pancreas with GAD65 antibody (clone GAD2/1960). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using GAD65 antibody (clone GAD2/1960). These results demonstrate the foremost specificity of the GAD2/1960 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free GAD65 antibody (clone GAD2/1960) as confirmation of integrity and purity.

Description

This mAb recognizes a protein of 65kDa, which is identified as glutamic acid decarboxylase 2 (GAD2). It is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. There are two forms of glutamic acid decarboxylases (GADs) that are found in the brain: GAD2 (also known as GAD65) and GAD1 (also known as GAD67). GAD1 and GAD2 are members of the group II decarboxylase family of proteins and are responsible for catalyzing the rate-limiting step in the production of GABA (gamma-aminobutyric acid) from L-glutamic acid. Although both GADs are found in the brain, GAD2 localizes to synaptic vesicle membranes in nerve terminals, while GAD1 is distributed throughout the cell. A pathogenic role for GAD2 is identified in the human pancreas since it has been identified as an autoantibody and an auto-reactive T cell target in insulin-dependent diabetes.

For highly specific detection of GAD65 in inhibitory synaptic signaling studies, see our [GAD65 Antibody / Synaptic GABA Marker Antibody](#) page featuring clone GAD2/2362 with strong HuProt(TM) microarray specificity validation data.

Application Notes

Optimal dilution of the GAD65 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 6-99 from the human protein were used as the immunogen for this GAD65 antibody.

Storage

Store the GAD65 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

