


## GADD45A Antibody (F54774)

| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F54774-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F54774-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

**Bulk quote request**


|                           |  |
|---------------------------|--|
| <b>Availability</b>       | 1-3 business days  |
| <b>Species Reactivity</b> | Human  |
| <b>Format</b>             | Purified   |
| <b>Clonality</b>          | Polyclonal (rabbit origin)   |
| <b>Isotype</b>            | Rabbit Ig  |
| <b>Purity</b>             | Purified   |
| <b>UniProt</b>            | P24522   |
| <b>Applications</b>       | Western Blot : 1:500-1:2000<br>Flow Cytometry : 1:25 (1x10 <sup>6</sup> cells)<br>Immunohistochemistry (FFPE) : 1:25 |
| <b>Limitations</b>        | This GADD45A antibody is available for research use only.  |




IHC testing of FFPE human tonsil tissue with GADD45A antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

kDa  
 130  
 72  
 55  
 36  
 28  
 17

Western blot testing of Y79 cell lysate with GADD45A antibody. Expected molecular weight: 14-18 kDa (monomer) and 28-36 kDa (dimer).



Western blot testing of 1) non-transfected and 2) transfected 293 cell lysate with GADD45A antibody. Expected molecular weight: 14-18 kDa (monomer) and 28-36 kDa (dimer).



Flow cytometry testing of human K562 cells with GADD45A antibody; Blue=isotype control, Green= GADD45A antibody.

## Description

GADD45A responds to environmental stresses by mediating activation of the p38/JNK pathway via MTK1/MEKK4 kinase. The GADD45A gene is a member of a group of genes whose transcript levels are increased following stressful growth arrest conditions and treatment with DNA-damaging agents. The DNA damage-induced transcription of this gene is mediated by both p53-dependent and -independent mechanisms.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the GADD45A antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 1-30 from the human protein was used as the immunogen for the GADD45A antibody.

## Storage

Aliquot the GADD45A antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.