


Galectin 3 Antibody (R31507)


Catalog No.	Formulation	Size
R31507	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
Gene ID	3958
Localization	Cytoplasmic, cell surface
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-4ug/ml Immunofluorescence (FFPE) : 5ug/ml Flow Cytometry : 1-3ug/10 ⁶ cells
Limitations	This Galectin 3 antibody is available for research use only.




Western blot testing of human 1) HeLa, 2) SiHa, 3) U-87 MG, 4) RT4, 5) HaCaT, 6) PC-3 and 7) T-47D cell lysate with Galectin 3 antibody. Predicted molecular weight ~26 kDa.




IHC-P: Galectin 3 antibody testing of human lung cancer tissue. HIER: steam section in pH6 citrate buffer for 20 min.




IHC-P: Galectin 3 antibody testing of human placenta tissue. HIER: steam section in pH6 citrate buffer for 20 min.



Immunofluorescent staining of FFPE human U-2 OS cells with Galectin 3 antibody at 2ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.



Immunofluorescent staining of FFPE human SiHa cells with Galectin 3 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human U-2 OS cells with Galectin 3 antibody at 1ug/10⁶ cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Galectin 3 antibody.

Description

Galectin 3 (GAL3), also known as LGALS3, MAC2 or GALBP, is a member of the lectin family, of which 14 mammalian galectins have been identified. Galectin-3 is encoded by a single gene, it located on chromosome 14, locus q21-q22. It is expressed in the nucleus, cytoplasm, mitochondrion, cell surface, and extracellular space. Studies have also shown that the expression is implicated in a variety of processes associated with heart failure, including myofibroblast proliferation, fibrogenesis, tissue repair, inflammation, and Ventricular remodeling. Galectin 3 is expressed in various tissues and organs, but is significantly absent in normal hepatocytes.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Galectin 3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Human partial recombinant protein (AA 139-250) was used as the immunogen for this Galectin 3 antibody.

Storage

After reconstitution, the Galectin 3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.