


Gamma Parvin Antibody / PARVG [clone 8C5.2] (V9268)


Catalog No.	Formulation	Size
V9268-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9268-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9268SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	8C5.2
Purity	Protein A/G affinity
UniProt	Q9HBI0
Localization	Cytoplasm, Cell Surface
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Gamma Parvin antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with Gamma Parvin antibody (clone 8C5.2) at 2ug/ml in PBS. Negative control inset: PBS instead of primary antibody to control for secondary binding. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Gamma Parvin antibody (clone 8C5.2) as confirmation of integrity and purity.

Description

Gamma-parvin likely plays a role in the regulation of cell adhesion and cytoskeleton organization. [UniProt]

Application Notes

Optimal dilution of the Gamma Parvin antibody should be determined by the researcher.

Immunogen

GST-tagged recombinant protein corresponding to human Gamma-parvin was used as the immunogen for the Gamma Parvin antibody.

Storage

Aliquot the Gamma Parvin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.