
# Gamma Parvin Antibody / PARVG [clone 8C5.2] (V9268)


| Catalog No. | Formulation | Size |
|---|---|---|
| V9268-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V9268-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V9268SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Purified |
| **Host** | Mouse |
| **Clonality** | Monoclonal (mouse origin) |
| **Isotype** | Mouse IgG1, kappa |
| **Clone Name** | 8C5.2 |
| **Purity** | Protein A/G affinity |
| **UniProt** | Q9HBI0 |
| **Localization** | Cytoplasm, Cell Surface |
| **Applications** | Western blot: 1-2ug/ml<br>Immunohistochemistry (FFPE): 1-2ug/ml |
| **Limitations** | This Gamma Parvin antibody is available for research use only. |



IHC staining of FFPE human tonsil tissue with Gamma Parvin antibody (clone 8C5.2) at 2ug/ml in PBS. Negative control inset: PBS instead of primary antibody to control for secondary binding. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




SDS-PAGE analysis of purified, BSA-free Gamma Parvin antibody (clone 8C5.2) as confirmation of integrity and purity.

## Description

Gamma-parvin likely plays a role in the regulation of cell adhesion and cytoskeleton organization. [UniProt]

## Application Notes

Optimal dilution of the Gamma Parvin antibody should be determined by the researcher.

## Immunogen

GST-tagged recombinant protein corresponding to human Gamma-parvin was used as the immunogen for the Gamma Parvin antibody.

## Storage

Aliquot the Gamma Parvin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.