


## GFAP Antibody Cocktail [clone GA-5 + ASTRO/789] (V2529)


Catalog No.	Formulation	Size
V2529-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2529-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2529SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2529IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	GA-5 + ASTRO/789
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P14136
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 0.25-0.5ug/ml for 30 min at RT
<b>Limitations</b>	This GFAP antibody cocktail is available for research use only.




IHC: Formalin-fixed, paraffin-embedded human Schwannoma stained with GFAP antibody cocktail (GA-5 + ASTRO/789).



IHC: Formalin-fixed, paraffin-embedded rat cerebellum stained with GFAP antibody cocktail (GA-5 + ASTRO/789).



IHC: Formalin-fixed, paraffin-embedded human cerebellum stained with GFAP antibody cocktail (GA-5 + ASTRO/789).



Western blot testing of human brain lysate with GFAP antibody cocktail (clones GA-5 + ASTRO/789). Predicted molecular weight ~50 kDa.

## Description

This mAb recognizes a protein of ~50kDa which is identified as Glial Fibrillary Acidic Protein (GFAP). It shows no cross-reaction with other intermediate filament proteins. GFAP is specifically found in astroglia. GFAP is a very popular marker for localizing benign astrocyte and neoplastic cells of glial origin in the central nervous system. Antibody to GFAP is useful in differentiating primary gliomas from metastatic lesions in the brain and for documenting astrocytic differentiation in tumors outside the CNS.

## Application Notes

Optimal dilution of the GFAP antibody cocktail should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

GFAP isolated from pig spinal cord (GA-5) and recombinant protein (ASTRO/789) were used as the immunogen for the GFAP antibody cocktail.

## Storage

Store the GFAP antibody cocktail at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

