


## GLAST Antibody / SLC1A3 / EAAT1 (N-Terminal Region) (F54102)

Catalog No.	Formulation	Size
F54102-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F54102-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Predicted Reactivity</b>	Bovine
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	P43003
<b>Applications</b>	Western Blot : 1:8000
<b>Limitations</b>	This GLAST antibody is available for research use only.



Western blot testing of GLAST antibody at 1:8000: Lane 1) human brain, 2) mouse brain and 3) mouse cerebellum lysate. Predicted molecular weight: 55-60 kDa.

### Description

Glutamate/aspartate transporter 1 (GLAST1) transports L-glutamate and also L- and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium. [UniProt]

### Application Notes

The stated application concentrations are suggested starting points. Titration of the GLAST antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## **Immunogen**

A portion of amino acids 127-161 from human GLAST was used as the immunogen for the GLAST antibody.

## **Storage**

Aliquot the GLAST antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.